



SHORT REPORT

Open Access

Confusions in orbivirus protein classification

Meik Dilcher* and Manfred Weidmann

Abstract

An extensive comparative analysis of orbivirus genomes revealed four cases of unclear numeration and protein designation, due to confused reference to protein size or segment size by which they are encoded. A concise nomenclature based on type species, sequence homology and functional characteristics independent of segment or protein size is suggested.

Keywords: *Reoviridae*, *Orbivirus*, Protein classification

Background

The genus *Orbivirus* is one of 15 in the family *Reoviridae* containing 22 serogroups (species) and at least 160 different serotypes (strains) [1]. Orbiviruses are transmitted by insects (midges, flies, mosquitoes) or by ticks. Their double-stranded RNA (dsRNA) genomes consist of 10 segments coding for seven structural and at least three non-structural proteins. Orbiviruses have no envelope but a double-shelled icosahedral capsid [2] and include pathogenic agents of wild animals (Epizootic hemorrhagic disease virus (EHDV)), domestic animals (Bluetongue virus (BTV) and African horse sickness virus (AHSV)), and of man (Kemerovo virus (KEMV)) [3]. Type species of the genus is the *Culicoides* midge transmitted BTV. Insect-borne orbiviruses are much better characterized than tick-transmitted orbiviruses for which few sequences have been described: Broadhaven virus (BRDV, partial) [4], Sandy Bay virus (SBaV, partial (formerly Nugget virus)) [5-8], St Croix River virus (SCRV, complete genome) [9], Great Island virus (GIV, complete genome) [7].

Recently we determined the complete genomes of Tri-beč virus (TRBV) and KEMV in a pyrosequencing approach [10] complementing available partial information on segments 1, 2 and 6 of these viruses and of Lipovnik virus (LIPV) [7].

During our extensive comparative analysis of orbivirus genomes we noticed four cases of unclear numeration and protein designation (see Tables 1 and 2). Some laboratories classify orbivirus proteins according to the

size of the proteins whereas others use the size of genome segments from which they are encoded.

- (i) The inner shell protein T2 for example can be encoded by segment 2 (tick- and mosquito-borne orbiviruses) or segment 3 (*Culicoides*-borne orbiviruses). This leads to some laboratories labeling this protein VP2(T2) (e.g. GIV), while others designate it VP3(T2) as in the type species BTV. For Peruvian horse sickness virus (PHSV) however, the segment 2 encoded protein is designated VP3 (T2) although it is larger (925 amino acids) than the segment 3 encoded VP2 protein (881 amino acids) [11]. To avoid confusion with the outer shell protein VP2 we suggest to exclusively use VP3(T2) for all T2 proteins.
- (ii) VP2 and VP2 homologous proteins can be encoded by segments 2, 3, 4 and 5 and are designated VP2, VP3 (YUOV, SCRIV) or VP4 (BRDV segment 4 ([12,13], sequence entry to GenBank missing)), GIV segment 5). Because of the location on the outer capsid and the described sequence similarity with other VP2 proteins, we suggest that the VP4 proteins (BRDV, GIV) as well as the VP3 proteins (YUOV, SCRIV) should be uniformly termed VP2, even though tick-borne VP2 proteins have only half the size of insect-borne VP2 proteins [13].
- (iii) The capping enzyme VP4(CaP) can be encoded by segment 4 (BTV, YUOV, SCRIV etc.) or segment 3 (TRBV, KEMV). In GIV this protein is designated VP3(CaP) [7] and should be renamed VP4(CaP) to avoid confusions with VP3(T2).
- (iv) In most cases VP5 is encoded by segment 6 and comprises a component of the outer shell that

* Correspondence: meik.dilcher@medizin.uni-goettingen.de
Department of Virology, University Medical Center Göttingen, Kreuzberggring 57, D-37075 Göttingen, Germany

Table 1 Comparison of the genome segments and encoded proteins of BTV, YUOV, TRBV, KEMV and GIV

BTV (insect-transmitted)				YUOV (insect-transmitted)				TRBV (tick-transmitted)			
Segment 1	VP1 (Pol)	150 kDa	RNA-dep.-RNA-Polymerase	Segment 1	VP1 (Pol)	151 kDa	RNA-dep.-RNA-Polymerase	Segment 1	VP1 (Pol)	146 kDa	RNA-dep.-RNA-Polymerase
3944 bp	ACR58458	1302 AA		3393 bp	YP_443925	1315 AA		3892 bp	HQ266581	1284 AA	
Segment 2	VP2	111 kDa	Outer shell	Segment 2	VP2 (T2)	107 kDa	Inner shell	Segment 2	VP3 (T2)	102 kDa	Inner shell
2953 bp	ACR58459	956 AA		2900 bp	YP_443926	940 AA		2793 bp	HQ266582	908 AA	
Segment 3	VP3 (T2)	103 kDa	Inner shell	Segment 3	VP3	100 kDa	Outer shell	Segment 3	VP4 (CaP)	72 kDa	Capping Enzyme
2772 bp	ACR58460	901 AA		2688 bp	YP_443927	873 AA		1935 bp	HQ266583	628 AA	
Segment 4	VP4 (CaP)	75 kDa	Capping Enzyme	Segment 4	VP4 (CaP)	74 kDa	Capping Enzyme	Segment 4	NS1 (TuP)	62 kDa	Formes Tubules
1980 bp	ACR58461	644 AA		1993 bp	YP_443928	645 AA		1734 bp	HQ266584	529 AA	
Segment 5	NS1 (TuP)	64 kDa	Forms Tubules	Segment 5	NS1 (TuP)	67 kDa	Forms Tubules	Segment 5	VP2	62 kDa	Outer shell
1769 bp	ACR58463	552 AA		1957 bp	YP_443929	574 AA		1730 bp	HQ266585	554 AA	
Segment 6	VP5	59 kDa	Outer shell	Segment 6	VP5	59 kDa	Outer shell	Segment 6	VP5	59 kDa	Outer shell
1638 bp	ACR58462	526 AA		1683 bp	YP_443930	535 AA		1668 bp	HQ266586	537 AA	
Segment 7	VP7 (T13)	39 kDa	Inner shell	Segment 7	NS2 (ViP)	48 kDa	Viral inclusion body matrix protein	Segment 7	NS2 (ViP)	41 kDa	Viral inclusion body matrix protein
1156 bp	ACR58464	349 AA		1504 bp	YP_443931	435 AA		1196 bp	HQ266587	368 AA	
Segment 8	NS2 (ViP)	41 kDa	Viral inclusion body matrix protein	Segment 8	VP7 (T13)	40 kDa	Inner shell	Segment 8	VP7 (T13)	40 kDa	Inner shell
1125 bp	ACR58465	354 AA		1191 bp	YP_443932	355 AA		1184 bp	HQ266588	357 AA	
Segment 9	VP6 (Hel)	36 kDa	ssRNA and dsRNA binding helicase	Segment 9	VP6 (Hel)	37 kDa	ssRNA and dsRNA binding helicase	Segment 9	VP6 (Hel)	33 kDa	ssRNA and dsRNA binding helicase
1049 bp	ACR58466	329 AA		1082 bp	YP_443933	338 AA		1034 bp	HQ266589	312 AA	
Segment 10	NS3	26 kDa	Glycoprotein	Segment 10	NS3	28 kDa	Glycoprotein	Segment 10	NS3	23 kDa	Glycoprotein
822 bp	ACR58467	229 AA		825 bp	YP_443934	253 AA		705 bp	HQ266590	214 AA	

KEMV (tick-transmitted)				GIV (tick-transmitted)			
Segment 1	VP1 (Pol)	146 kDa	RNA-dep.-RNA-Polymerase	Segment 1	VP1 (Pol)	147 kDa	RNA-dep.-RNA-Polymerase
3896 bp	HQ266591	1285 AA		3897 bp	ADM88592	1285 AA	
Segment 2	VP3 (T2)	103 kDa	Inner shell	Segment 2	VP2 (T2)	103 kDa	Inner shell
2792 bp	HQ266592	908 AA		2794 bp	ADM88593	908 AA	
Segment 3	VP4 (CaP)	72 kDa	Capping Enzyme	Segment 3	VP3 (CaP)	73 kDa	Capping Enzyme
1934 bp	HQ266593	632 AA		1936 bp	ADM88594	635 AA	
Segment 4	VP2	63 kDa	Outer shell	Segment 4	NS1 (TuP)	60 kDa	Formes Tubules
1730 bp	HQ266594	554 AA		1731 bp	ADM88595	531 AA	
Segment 5	NS1 (TuP)	60 kDa	Formes Tubules	Segment 5	VP4	62 kDa	Outer shell
1719 bp	HQ266595	529 AA		1722 bp	ADM88596	551 AA	
Segment 6	VP5	59 kDa	Outer shell	Segment 6	VP5	60 kDa	Outer shell
1668 bp	HQ266596	537 AA		1666 bp	ADM88597	537 AA	
Segment 7	NS2 (ViP)	41 kDa	Viral inclusion body matrix protein	Segment 7	VP7 (T13)	40 kDa	Inner shell
1197 bp	HQ266597	368 AA		1181 bp	ADM88598	357 AA	
Segment 8	VP7 (T13)	40 kDa	Inner shell	Segment 8	NS2 (ViP)	39 kDa	Viral inclusion body matrix protein
1183 bp	HQ266598	357 AA		1172 bp	ADM88599	359 AA	
Segment 9	VP6 (Hel)	34 kDa	ssRNA and dsRNA binding helicase	Segment 9	VP6 (Hel)	34 kDa	ssRNA and dsRNA binding helicase
1049 bp	HQ266599	317 AA		1056 bp	AMD88600	321 AA	
Segment 10	NS3	23 kDa	Glycoprotein	Segment 10	NS3	19 kDa	Glycoprotein
707 bp	HQ266600	214AA		703 bp	ADM88602	171 AA	

Outer and inner shell proteins are labeled in bold. GenBank and SwissProt accession numbers are indicated.

Table 2 Comparison of the genome segments and encoded proteins of SCRV, PHSV, BRDV and LIPV

SCRV (tick-transmitted)				PHSV (isolates only known from horses)			
Segment 1	VP1 (Pol)	151 kDa	RNA-dep.-RNA-Polymerase	Segment 1	VP1 (Pol)	151 kDa	RNA-dep.-RNA-Polymerase
4089 bp	YP_052942	1345 AA		3987 bp	YP_460038	1311 AA	
Segment 2	VP2 (T2)	98 kDa	Inner shell	Segment 2	VP3 (T2)	105 kDa	Inner shell
2747 bp	YP_052943	890 AA		2856 bp	YP_460039	925 AA	
Segment 3	VP3	74 kDa	Outer shell	Segment 3	VP2	104 kDa	Outer shell
2024 bp	YP_052944	654 AA		2747 bp	YP_460040	881 AA	
Segment 4	VP4 (CaP)	74 kDa	Capping Enzyme	Segment 4	VP4 (CaP)	74 kDa	Capping Enzyme
2017 bp	YP_052945	643 AA		1996 bp	YP_460041	646 AA	
Segment 5	VP5	57 kDa	Outer shell	Segment 5	NS1 (TuP)	64 kDa	Forms Tubules
1664 bp	YP_052946	517 AA		1784 bp	YP_460045	554 AA	
Segment 6	NS1 (TuP)	58 kDa	Forms Tubules	Segment 6	VP5	59 kDa	Outer shell
1657 bp	YP_052947	517 AA		1695 bp	YP_460042	529 AA	
Segment 7	NS2 (ViP)	51 kDa	Viral inclusion body matrix protein	Segment 7	NS2 (ViP)	48 kDa	Viral inclusion body matrix protein
1463 bp	YP_052948	462 AA		1613 bp	YP_460046	435 AA	
Segment 8	VP7 (T13)	41 kDa	Inner shell	Segment 8	VP7 (T13)	40 kDa	Inner shell
1256 bp	YP_052949	379 AA		1180 bp	YP_460044	353 AA	
Segment 9	VP6 (Hel)	26 kDa	ssRNA and dsRNA binding helicase	Segment 9	VP6 (Hel)	37 kDa	ssRNA and dsRNA binding helicase
764 bp	YP_052950	232 AA		1071 bp	YP_460043	334 AA	
Segment 10	NS3	24 kDa	Glycoprotein	Segment 10	NS3	28 kDa	Glycoprotein
764 bp	YP_052951	224 AA		819 bp	YP_460047	255 AA	

BRDV (tick-transmitted)				LIPV (tick-transmitted)			
Segment 1				Segment 1	VP1 (Pol)	146 kDa	RNA-dep.-RNA-Polymerase
				3892 bp	ADM88603	1284 AA	
Segment 2	VP2 (T2)	103 kDa	Inner shell	Segment 2	VP2 (T2)	103 kDa	Inner shell
	P35934	908 AA		2793 bp	ADM88604	908 AA	
Segment 3				Segment 3			
Segment 4	VP4 †	63 kDa †	Outer shell	Segment 4			
Segment 5	VP5	53 kDa	Outer shell	Segment 5			
1658 bp	P21230	480 AA					
Segment 6	NS1 (TuP)	60 kDa	Formes Tubules	Segment 6	VP5	502 AA ††	Outer shell
1714 bp	2115436A	537 AA		1509 bp ††	ADM88605		
Segment 7	VP7 (T13)	40 kDa	Inner shell	Segment 7			
	P35935	356 AA					
Segment 8				Segment 8			
Segment 9				Segment 9			
Segment 10	NS3	22 kDa	Glycoprotein	Segment 10			
	P32555	205 AA					

Outer and inner shell proteins are labeled in bold. GenBank and SwissProt accession numbers are indicated.

†: [12,13], GenBank entry missing.

††: partial sequence.

might be involved in membrane fusion and penetration [14]. TRBV and KEMV also encode VP5 on segment 6. The highest similarity of TRBV VP5 is to LIPV VP5 (95.6%), again encoded by segment 6 [7]. However, VP5 of BRDV is described as encoded by segment 5 [15]. Since in the classification of the viral genome segments bigger segments have smaller segment numbers, and the size of BRDV segment 6 (1714 bp) encoding the NS1(TuP) [16] is larger than the size of BRDV segment 5 (1658 bp) encoding VP5, a reassignment of BRDV segment 5 and 6 (a vice versa switch) seems necessary.

To summarize, it would be much more helpful if the nomenclature of the viral proteins in orbiviruses would reflect the sequence homology and functional relationship rather than protein size or encoding segment size, since the sizes of the orbivirus genome segments sometimes only differ slightly, which leads to even closely related viruses such as TRBV and KEMV encoding VP2 and NS1 (TuP) on different genome segments. We therefore suggest the following concise nomenclature based on the type species BTV and on sequence homology and functional characteristics independent of segment or protein size: VP1(Pol), VP2, VP3(T2), VP4(CaP), VP5, VP6(Hel), VP7 (T13), NS1(TuP), NS2(ViP), NS3.

Competing interests

The authors declare that they have no competing interests.

Author's contributions

MD and MW wrote the paper. Both authors read and approved the final manuscript.

Acknowledgments

This work was supported by the Federal Ministry of Education and Research (BMBF), grant number 01KI0710, "Research on Zoonotic Infectious Diseases" program, "Emerging arthropod-borne viral infections in Germany: Pathogenesis, diagnostics and surveillance" and the BMBF funded research program "Potential release-oriented biothreat emergency diagnostics (P.R.O.B. E)" for civil security of the German Federal Government as part of the high-tech strategy for Germany.

Received: 1 February 2012 Accepted: 24 July 2012

Published: 21 August 2012

References

1. Attoui H, Mertens PPC, Becnel J, Belagahanalli S, Bergoin M, Brussaard CP, Chappell JD, Ciarlet M, del Vas M, Dermody TS, et al: **Orbiviruses, Reoviridae**. In *Virus taxonomy ninth report of the international committee on taxonomy of viruses*. Edited by King AMQ, Adams MJ, Carstens EB, Lefkowitz EJ. Elsevier: Academic Press; 2011:592–603.
2. Attoui H, Jaafar FM, Belhouchet M, Aldrovandi N, Tao SJ, Chen BQ, Liang GD, Tesh RB, de Micco P, de Lamballerie X: **Yunnan orbivirus, a new orbivirus species isolated from Culex tritaeniorhynchus mosquitoes in China**. *J Gen Virol* 2005, **86**:3409–3417.
3. Gorman BM: **Variation in orbiviruses**. *J Gen Virol* 1979, **44**:1–15.
4. Moss SR, Jones LD, Nuttall PA: **Comparison of the major structural core proteins of tick-borne and culicoides-borne orbiviruses**. *J Gen Virol* 1992, **73**:2585–2590.
5. Doherty RL, Carley JG, Murray MD, Main AJ, Kay BH, Domrow R: **Isolation of arboviruses (Kemerovo-Group, Sakhalin-Group) from ixodes-uriae collected at Macquarie Island, Southern Ocean**. *Am J Trop Med Hyg* 1975, **24**:521–526.
6. Gorman BM, Taylor J, Morton HC, Melzer AJ, Young PR: **Characterization of nugget virus, a serotype of the kemeroovo group of orbiviruses**. *Aust J Exp Biol Med Sci* 1984, **62**:101–115.
7. Belhouchet M, Jaafar FM, Tesh RB, Grimes J, Maan S, Mertens P, Attoui H: **Complete sequence of the Great Island virus and comparison with the T2 and outer-capsid proteins of Kemerovo, Lipovnik and Tribec viruses (genus Orbivirus, family Reoviridae)**. *J Gen Virol* 2010, **91**:2985–2993.
8. Major L, Linn ML, Slade RW, Schroder WA, Hyatt AD, et al: **Ticks associated with Macquarie Island penguins carry arboviruses from four genera**. *PLoS One* 2009, **4**:e4375.
9. Attoui H, Stirling JM, Munderloh UG, Billoir F, Brookes SM, Burroughs JN, de Micco P, Mertens PPC, de Lamballerie X: **Complete sequence characterization of the genome of the St Croix River virus, a new orbivirus isolated from cells of Ixodes scapularis**. *J Gen Virol* 2001, **82**:795–804.
10. Dilcher M, Hasib L, Lechner M, Wieseke N, Middendorf M, Marz M, Koch A, Spiegel M, Dobler G, Hufert FT, Weidmann M: **Genetic characterization of Tribec virus and Kemerovo virus, two tick-transmitted human-pathogenic orbiviruses**. *Virology* 2012, **423**:68–76.
11. Attoui H, Mendez-Lopez MR, Rao SJ, Hurtado-Alendes A, Lizaraso-Caparo F, Jaafar FM, Samuel AR, Belhouchet M, Pritchard LI, Melville L, et al: **Peruvian horse sickness virus and Yunnan orbivirus, isolated from vertebrates and mosquitoes in Peru and Australia**. *Virology* 2009, **394**:298–310.
12. Moss SR, Nuttall PA: **Subcore-like and core-like particles of Broadhaven Virus (Brdv), a tick-borne orbivirus, synthesized from baculovirus-expressed Vp2 and Vp7, the major core proteins of Brdv**. *Virus Res* 1994, **32**:401–407.
13. Schoehn G, Moss SR, Nuttall PA, Hewat EA: **Structure of Broadhaven virus by cryoelectron microscopy: Correlation of structural and antigenic properties of Broadhaven virus and bluetongue virus outer capsid proteins**. *Virology* 1997, **235**:191–200.
14. Zhang X, Boyce M, Bhattacharya B, Zhang XK, Schein S, Roy P, Zhou ZH: **Bluetongue virus coat protein VP2 contains sialic acid-binding domains, and VP5 resembles enveloped virus fusion proteins**. *Proc Natl Acad Sci U S A* 2010, **107**:6292–6297.
15. Moss SR, Fukusho A, Nuttall PA: **Rna segment 5 of Broadhaven virus, a tick-borne orbivirus, shows sequence homology with segment-5 of bluetongue virus**. *Virology* 1990, **179**:482–484.
16. Moss SR, Nuttall PA: **Comparison of the nonstructural protein, Ns1, of tick-borne and insect-borne orbiviruses**. *Virus Res* 1995, **36**:287–292.

doi:10.1186/1743-422X-9-166

Cite this article as: Dilcher and Weidmann: Confusions in orbivirus protein classification. *Virology Journal* 2012 **9**:166.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

